

Volume contents

Volume 74 (1998)

| | | | |
|---|-----|---|-----|
| Issue contents | 1 | related to the <i>Ski</i> proto-oncogene and are expressed in eye and limb | 121 |
| N.E. Baker, S.-Y. Yu | | G. Pillemer, R. Yelin, M. Epstein, L. Gont, Y. Frumkin, J.K. Yisraeli, H. Steinbeisser, A. Fainsod | |
| The R8-photoreceptor equivalence group in <i>Drosophila</i> : fate choice precedes regulated <i>Delta</i> transcription and is independent of <i>Notch</i> gene dose | 3 | The <i>Xcad-2</i> gene can provide a ventral signal independent of <i>BMP-4</i> | 133 |
| B.K. Mueller, D. Dütting, A. Haase, A. Feucht, P. Macchi | | K. Itoh, J. Jacob, S.Y. Sokol | |
| Partial respecification of nasotemporal polarity in double-temporal chick and chimeric chick-quail eyes | 15 | A role for <i>Xenopus</i> Frizzled 8 in dorsal development | 145 |
| A. Metz, S. Knöchel, P. Büchler, M. Köster, W. Knöchel | | Gene expression patterns | |
| Structural and functional analysis of the <i>BMP-4</i> promoter in early embryos of <i>Xenopus laevis</i> | 29 | F. Loosli, R.W. Köster, M. Carl, A. Krone, J. Wittbrodt | |
| S.M. Bell, C.M. Schreiner, W.J. Scott | | <i>Six3</i> , a medaka homologue of the <i>Drosophila</i> homeobox gene <i>sine oculis</i> is expressed in the anterior embryonic shield and the developing eye | 159 |
| The loss of ventral ectoderm identity correlates with the inability to form an AER in the <i>legless</i> hindlimb bud | 41 | J.J. Gibson-Brown, S.I. Agulnik, L.M. Silver, V.E. Papaioannou | |
| M. Fanto, C.A. Mayes, M. Mlodzik | | Expression of T-box genes <i>Tbx2-Tbx5</i> during chick organogenesis | 165 |
| Linking cell-fate specification to planar polarity: determination of the R3/R4 photoreceptors is a prerequisite for the interpretation of the Frizzled mediated polarity signal | 51 | S. Nayak, N. Galili, C.A. Buck | |
| N. Kahane, Y. Cinnamon, C. Kalcheim | | Immunohistochemical analysis of the expression of two serine-threonine kinases in the maturing mouse testis | 171 |
| The origin and fate of pioneer myotomal cells in the avian embryo | 59 | Y. Maruoka, N. Ohbayashi, M. Hoshikawa, N. Itoh, B.L.M. Hogan, Y. Furuta | |
| S.-i. Nishimatsu, G.H. Thomsen | | Comparison of the expression of three highly related genes, <i>Fgf8</i> , <i>Fgf17</i> and <i>Fgf18</i> , in the mouse embryo | 175 |
| Ventral mesoderm induction and patterning by bone morphogenetic protein heterodimers in <i>Xenopus</i> embryos | 75 | T. Jaskoll, H. Chen, P.C. Denny, P.A. Denny, M. Melnick | |
| F. Levavasseur, W. Mandemakers, P. Visser, L. Broos, F. Grosveld, D. Zivkovic, D. Meijer | | Mouse submandibular gland mucin: embryo-specific mRNA and protein species | 179 |
| Comparison of sequence and function of the <i>Oct-6</i> genes in zebrafish, chicken and mouse | 89 | D. Bellomonte, M. Di Bernardo, R. Russo, G. Caronia, G. Spinelli | |
| A. Prokop, S. Bray, E. Harrison, G.M. Technau | | Highly restricted expression at the ectoderm-endoderm boundary of <i>PIHbox 9</i> , a sea urchin homeobox gene related to the human <i>HB9</i> gene | 185 |
| Homeotic regulation of segment-specific differences in neuroblast numbers and proliferation in the <i>Drosophila</i> central nervous system | 99 | T. Léveillard, P. Gorrry, K. Niederreither, B. Wasylyk | |
| J.B.L. Bard, M.H. Kaufman, C. Dubreuil, R.M. Brune, A. Burger, R.A. Baldock, D.R. Davidson | | MDM2 expression during mouse embryogenesis and the requirement of p53 | 189 |
| An internet-accessible database of mouse developmental anatomy based on a systematic nomenclature | 111 | Author Index | 195 |
| K.L. Hammond, I.M. Hanson, A.G. Brown, L.A. Lettice, R.E. Hill | | Subject Index | 197 |
| Mammalian and <i>Drosophila dachshund</i> genes are | | Volume Contents | 205 |